

Have you ever had an ICU patient...

- ▶ become progressively more swollen and edematous after fluid resuscitation?
- ▶ develop renal failure and need dialysis?
- ▶ develop multiple organ failure and die?

What was their Intra-Abdominal Pressure?

AbViser® AutoValve® – IAP Monitoring Device is the first and only closed IAP monitoring system with proven accuracy, reproducibility and safety.¹

- **Simplifies ICU Nursing Tasks** –
Easy set-up and use
- **Standardized Data** –
Proven accuracy, reproducibility and safety.¹
- **Automatic Valve with Time Release** –
Eliminates clamping and turning of stopcocks
- **White Stripe Tubing**
(designates IAP) –
Reduces misconnection errors
- **Closed System w/Syringe Sleeve** –
Helps reduce nosocomial infection risk²
- **OR Compatible** –
Data assists surgeon in closing laparotomy
- **Early IAH Recognition** –
Consistent monitoring improves outcomes



Sepsis/SIRS + Volume Resuscitation = Intra-Abdominal Hypertension³⁻⁴




Monitoring IAP + Intervention = Reduced Organ Failure⁵⁻⁹



Elegant solutions for complex issues.

**AbViser®
AutoValve®**

Intra-Abdominal Pressure Monitoring Device



Ordering Information:

Reorder Number	Quantity / Case	Description
ABV300	10	AbViser [®] AutoValve [™] IAP Monitoring Device, Patient Mount
ABV301	10	AbViser [®] AutoValve [™] IAP Monitoring Device, Pole Mount
ABV320	10	AbViser [®] AutoValve [™] IAP Monitoring Device with Edwards Lifesciences TruWave [®] Transducer, Patient Mount
ABV321	10	AbViser [®] AutoValve [™] IAP Monitoring Device with Edwards Lifesciences TruWave [®] Transducer, Pole Mount
ABV330	10	AbViser [®] AutoValve [™] IAP Monitoring Device with Hospira Transpac [®] IV Transducer, Patient Mount
ABV331	10	AbViser [®] AutoValve [™] IAP Monitoring Device with Hospira Transpac [®] IV Transducer, Pole Mount
ABV350	5	AbViser [®] Neonate Adapter

[Latex Free] [Individually Packaged Sterile]

References:

- Kimball, E.J., et al., Reproducibility of bladder pressure measurements in critically ill patients. Intensive Care Med, 2007.
- Huey WY, Newton DW, Augustine SC, Verjaska BD, Mitrano FP. Microbial contamination potential of sterile disposable plastic syringes. American Journal of Hospital Pharmacy 1985; 42:102-5
- Cheatham, M.L., et al., Results from the International Conference of Experts on Intra-abdominal Hypertension and Abdominal Compartment Syndrome. II. Recommendations. Intensive Care Med, 2007.
- Malbrain, M.L.N.G., et al., Incidence and prognosis of intraabdominal hypertension in a mixed population of critically ill patients: a multiple-center epidemiological study. Crit Care Med, 2005. 33(2): p. 315-22.
- Joseph, D.K., et al., Decompressive laparotomy to treat intractable intracranial hypertension after traumatic brain injury. J Trauma, 2004. 57(4): p. 687-95.
- Sun, et al., Indwelling catheter and conservative measures in the treatment of abdominal compartment syndrome in fulminant acute pancreatitis. World J Gastroenterol, 2006. 12(31): p. 5068-70.
- Oda, S., et al., Management of Intra-abdominal Hypertension in Patients With Severe Acute Pancreatitis With Continuous Hemodiafiltration Using a Polymethyl Methacrylate Membrane Hemofilter. Ther Apher Dial, 2005. 9(4): p. 355-61.
- Ivatury, R.R., et al., Intra-abdominal hypertension after life-threatening penetrating abdominal trauma: prophylaxis, incidence, and clinical relevance to gastric mucosal pH and abdominal compartment syndrome. J Trauma, 1998. 44(6): p. 1016-21.
- Sugrue, M., et al., Intra-abdominal hypertension is an independent cause of postoperative renal impairment. Arch Surg, 1999. 134(10): p. 1082-5.